



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Mortuary statistics from July 6 to July 12, 1902, inclusive, for the port of Santiago de Cuba.*

Typhoid fever (abdominal typhus) .....	1
Fever, intermittent malarial .....	2
La grippe .....	1
Tubercle of lungs.....	5
Cancer (organs not specified).....	1
Cerebral congestion.....	1
Tetanus.....	2
Bronchitis, acute.....	1
Diarrhea and enteritis (under 2 years)....	4
Bright's disease.....	1
Ill-defined or unspecified causes of death.....	2

Total..... 21

Annual rate of mortality for the week, 25.39 per 1,000. Estimated population, 43,000.

*Mortuary statistics from July 13 to July 19, 1902, inclusive, for the Port of Santiago de Cuba.*

Malarial cachexia.....	1
Diphtheritic paralysis .....	1
Dysentery .....	1
Tubercle of lungs.....	2
Meningitis .....	1
Organic diseases of the heart.....	2
Diarrhea and enteritis (under 2 years)....	2
Diarrhea and enteritis, chronic.....	1
Diarrhea and enteritis (2 years and over)....	2
Bright's disease.....	1
Septicæmia, puerperal.....	1
Congenital debility .....	1
Ill-defined or unspecified causes of death.....	1

Total ..... 17

Annual rate of mortality for the week, 20.5 per 1,000. Estimated population, 43,000.

## ENGLAND.

*Report from Liverpool.*

LIVERPOOL, ENGLAND, *July 22, 1902.*

SIR: I have the honor to state that for the week ended July 19, 1902, there were reported to the local health authorities the following cases: Smallpox, 9; typhus fever, 2; enteric fever, 15; scarlet fever, 72; diphtheria, 12; measles, 7. This is an increase in the number of cases of smallpox over the preceding weeks, and is probably due to fresh importations from London. For the same period in Birkenhead there were reported the following cases: Smallpox, 1; scarlet fever, 1; diphtheria, 2; membranous croup, 1; enteric fever, 4, and measles, 48. Between the 15th and 18th of this month there have been in Bootle 9 cases of smallpox, with 1 death in an unvaccinated person. Bootle is a corporation immediately adjacent to Liverpool, in fact, virtually a part of it.

Respectfully,

CARROLL FOX,  
*Assistant Surgeon.*

The SURGEON-GENERAL.